

UC Irvine

UC Irvine Previously Published Works

Title

Influence of biomass combustion emissions on the distribution of acidic trace gases over the southern Pacific basin during austral springtime

Permalink

<https://escholarship.org/uc/item/6bk523r9>

Journal

Journal of Geophysical Research Atmospheres, 104(D5)

ISSN

0148-0227

Authors

Talbot, RW
Dibb, JE
Scheuer, EM
et al.

Publication Date

1999-03-20

DOI

10.1029/98JD00879

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Influence of biomass combustion emissions on the distribution of acidic trace gases over the southern Pacific basin during austral springtime

R. W. Talbot,¹ J. E. Dibb,¹ E. M. Scheuer,¹ D. R. Blake,² N. J. Blake,² G. L. Gregory,³
G. W. Sachse,³ J. D. Bradshaw,^{4,5} S. T. Sandholm,⁴ and H. B. Singh⁶

Abstract. This paper describes the large-scale distributions of HNO_3 , HCOOH , and CH_3COOH over the central and South Pacific basins during the Pacific Exploratory Mission-Tropics (PEM-Tropics) in austral springtime. Because of the remoteness of this region from continental areas, low part per trillion by volume (pptv) mixing ratios of acidic gases were anticipated to be pervasive over the South Pacific basin. However, at altitudes of 2–12 km over the South Pacific, air parcels were encountered frequently with significantly enhanced mixing ratios (up to 1200 pptv) of acidic gases. Most of these air parcels were centered in the 3–7 km altitude range and occurred within the 15° – 65° S latitudinal band. The acidic gases exhibited an overall general correlation with CH_2Cl , PAN, and O_3 , suggestive of photochemical and biomass burning sources. There was no correlation or trend of acidic gases with common industrial tracer compounds (e.g., C_2Cl_4 or CH_3CCl_3). The combustion emissions sampled over the South Pacific basin were relatively aged exhibiting $\text{C}_2\text{H}_2/\text{CO}$ ratios in the range of 0.2–2.2 pptv/ppbv. The relationships between acidic gases and this ratio were similar to what was observed in aged air parcels (i.e., >3–5 days since they were over a continental area) over the western North Pacific during the Pacific Exploratory Mission-West Phases A and B (PEM-West A and B). In the South Pacific marine boundary layer a median $\text{C}_2\text{H}_2/\text{CO}$ ratio of 0.6 suggested that this region was generally not influenced by direct inputs of biomass combustion emissions. Here we observed the lowest mixing ratios of acidic gases, with median values of 14 pptv for HNO_3 , 19 pptv for HCOOH , and 18 pptv for CH_3COOH . These values were coincident with low mixing ratios of NO_x (<10 pptv), CO (\approx 50 parts per billion by volume (ppbv)), O_3 (< 20 ppbv), and long-lived hydrocarbons (e.g., C_2H_6 < 300 pptv). Overall, the PEM-Tropics data suggest an important influence of aged biomass combustion emissions on the distributions of acidic gases over the South Pacific basin in austral springtime.

1. Introduction

Acidic gases are important participants in tropospheric chemical processes. They are major end products of oxidative cycles, with wet and dry removal of HNO_3 and H_2SO_4 from the atmosphere principal sinks for tropospheric NO_x ($\text{NO} + \text{NO}_2$) and SO_2 [Logan, 1983; Hales and Dana, 1979]. In remote regions the monocarboxylic acids HCOOH and CH_3COOH are often the dominant acidic gases and acidity components of cloud water and precipitation [Keene *et al.*, 1983; Andreae *et al.*, 1988, 1990]. Formic acid is also a major sink for OH radicals in cloudwater [Jacob, 1986].

Formic acid may be produced by aqueous phase OH oxidation of hydrated formaldehyde ($\text{H}_2\text{C}(\text{OH})_2$) in cloudwater and subse-

quently provides an important source of gas phase HCOOH in the remote troposphere [Chameides and Davis, 1983; Jacob, 1986]. Aqueous phase production mechanisms for CH_3COOH appear to be quite slow and probably are a negligible source of this species to the troposphere [Jacob and Wofsy, 1988]. The major sources of HCOOH and CH_3COOH to the global troposphere appear to be emissions from combustion [Kawamura *et al.*, 1985; Talbot *et al.*, 1988; Helas *et al.*, 1992; Lefer *et al.*, 1994], vegetation [Keene and Galloway, 1986; Talbot *et al.*, 1988, 1990], and possibly soils [Sanhueza and Andreae, 1991; Talbot *et al.*, 1995]. Permutation reactions of peroxy radicals have been proposed as potentially important sources of carboxylic acids [Madronich and Calvert, 1990; Madronich *et al.*, 1990], but recent measurements at a continental site indicate that this pathway may be relatively unimportant [Talbot *et al.*, 1995].

There are potentially numerous production mechanisms for HNO_3 in the troposphere including, $\text{NO}_2 + \text{OH}$, recycling of reactive nitrogen reservoir species, and evaporation of NO_3^- in aerosol and aqueous phases [Roberts, 1995]. Many of these processes are thought to very slow in the upper tropical troposphere due to low O_3 mixing ratios and cold temperatures retaining most of the reactive nitrogen in the form of NO during the daytime [Folkins *et al.*, 1995].

Measurements conducted in winter 1992 at 10–12 km altitude between Tahiti and California showed an abrupt decrease in the mixing ratios of O_3 and NO_y (i.e., the sum of reactive nitrogen species) at the southern edge of the Intertropical Convergence Zone

¹Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham.

²Department of Chemistry, University of California, Irvine.

³NASA Langley Research Center, Hampton, Virginia.

⁴Department of Earth and Atmospheric Science, Georgia Institute of Technology, Atlanta.

⁵Deceased June 16, 1997.

⁶NASA Ames Research Center, Moffett Field, California.

(ITCZ) [Folkins *et al.*, 1995]. Owing to the remoteness of the South Pacific basin from continental areas, the observed trend in O_3 and NO_y is not surprising. In fact, low mixing ratios would be expected to be pervasive over the South Pacific basin for most tropospheric trace species, including acidic gases.

In this paper we present the large-scale distributions of HNO_3 , HCOOH , and CH_3COOH over the central and South Pacific basins during the NASA Global Tropospheric Experiment/Pacific Exploratory Mission-Tropics (GTE/PEM-Tropics) in September/October 1996. Objectives of PEM-Tropics included obtaining baseline data for important tropospheric gases, evaluating the oxidizing capacity of the troposphere and factors influencing it, and improving our understanding of the natural sulfur cycle over the South Pacific basin.

The first part of this paper focuses on the distributions of acidic gases in the 2–12 km altitude range which were apparently heavily impacted by aged biomass combustion emissions. Supporting evidence for this source is provided by coincident distributions of selected hydrocarbon compounds. The distribution of acidic gases is examined in the marine boundary layer and overlying transition region in the second part of this paper. Here there was little evidence for a direct influence of biomass combustion inputs on the chemistry, in stark contrast to the middle and upper troposphere. Overall, the PEM-Tropics measurements provide unique information of the chemistry of this extensive remote region during austral springtime.

2. Experimental Methods

2.1. Study Area

The PEM-Tropics airborne expedition was conducted using the NASA Ames DC-8 research aircraft. Transit and intensive site science missions composed 18 flights, averaging 8–10 hours in duration and covering the altitude range of 0.3 to 12.5 km. The base of operations for these missions progressed as follows: (1) Tahiti (three missions), (2) Easter Island (two missions), (3) Tahiti (one mission), (4) New Zealand (one mission), and (5) Fiji (three missions). The data used in this paper were obtained in the geographic grid approximately bounded by 60°N–75°S latitude and 165°E–105°W longitude. Data obtained on transit flights (eight missions) were also utilized in this paper. A geographic map of the study region is shown in several companion papers [e.g., Gregory *et al.*, this issue; Hoell *et al.*, this issue].

The overall scientific rationale and description of individual aircraft missions is described in the PEM-Tropics overview paper [Hoell *et al.*, this issue]. The features of the large-scale meteorological regime and associated air mass trajectory analyses for the September–October 1996 time period are presented by Fuelberg *et al.* [this issue].

2.2. Sampling and Analytical Methodology

Acidic gases were subsampled from a high-volume (500–1500 standard liters per minute (sLpm)) flow of ambient air using the mist chamber technique [Talbot *et al.*, 1988, 1990, 1997a]. The subsample flow rate was always <10% of the primary manifold total flow. Sample collection intervals were typically 4 min in the boundary layer, 6 min at 2–9 km altitude, and 8 min above 9 km altitude, reflecting decreased pumping rates in the middle and upper troposphere. The inlet manifold consisted of a 0.9 m length of 41 mm ID glass coated stainless steel pipe. The pipe extended from the DC-8 fuselage to provide a 90° orientation to the ambient air streamline flow. To facilitate pumping of the high-volume manifold

flow, a diffuser was mounted over the end of the inlet pipe parallel to the DC-8 fuselage. This device provided a "shroud" effect, slowing the flow of ambient air through it slightly below the true air speed of the DC-8 and adding 50–100 hPa of pressurization to the sampling manifold. This effectively eliminated the reverse venturi effect (≈ 40 hPa) on the sampling manifold. An additional feature of the diffuser was a curved step around the manifold pipe which provided the streamline effects of a backward facing inlet. Its function was to facilitate exclusion of aerosol particles greater than $\approx 2 \mu\text{m}$ in diameter from the sampling manifold. Aerosols smaller than this were removed from the sampled air stream using a 1 μm pore-sized Zeflur teflon filter that was readily changeable every 5–10 min to minimize aerosol loading on the filter and gas/aerosol phase partitioning from ambient conditions.

In addition to the features described above, the inlet manifold was equipped with the capability for conducting a standard addition of HNO_3 into the manifold ambient air stream. This spike was added ≈ 10 cm downstream inside the manifold pipe through a 6.5 mm OD glass coated stainless steel tube mounted perpendicular to the air flow. This tube was ≈ 20 mm long and maintained at 40°C to facilitate passing of calibration gas through it. Through a tee, this injection length of tubing was connected to about 1.5 m of heated tubing that was directly linked to the perm oven output. This design effectively tested the passing efficiency of the entire manifold system, which was indistinguishable from $100 \pm 15\%$. The calibration system for HNO_3 consisted of a permeation oven held at 50°C and a dilution flow of ultra zero air (1.5 sLpm) which swept the oven outflow to either a nylon filter for output quantification or the sampling manifold for standard addition on ambient air. The heated tubing through which the HNO_3 stream passed was kept equilibrated by a flow design that allowed the calibration gas to constantly pass to near the point of injection into the manifold flow before being dumped to waste through a return line. The mixing ratio of HNO_3 in the 1.5 sLpm flow was typically 200 parts per billion by volume (ppbv). This spike was then diluted several hundred times by the high flow rate of ambient air in the sampling manifold, producing standard additions of 100–1200 pptv. Previously we have studied the passing efficiency of carboxylic acids through our inlet manifold, and found it to be $\geq 95\%$ [Talbot *et al.*, 1992]. Thus we focused our attention on HNO_3 due to its importance to the reactive nitrogen cycle and tropospheric chemistry.

The permeation oven output of HNO_3 was monitored on the ground and in the air in near-real time. We fabricated a new calibration system for PEM-Tropics which maintained the permeation tube to $50.0 \pm 0.1^\circ\text{C}$ and 1850 ± 1 hPa pressure at all altitudes. The permeation source was constant to $\pm 8.5\%$ over the course of the expedition, with no equilibration time required at any altitude, even with rapid changes such as during spiral maneuvers. This new design utilized upstream pressure control (i.e., before the permeation tube) so that there were no fittings, valves, or flow/pressure controllers in-line between the tube and the injection point into our sampling manifold. The flow through the oven varied from 20–25 $\text{cm}^3 \text{min}^{-1}$ depending on the ambient pressure and was diluted into the 1.5 sLpm flow described above. Standard additions were conducted with and without a teflon filter in-line to verify that the filter did not influence the passing efficiency of the sampling manifold.

Computer controlled syringe pumps were used to move sample solutions in and out of the mist chamber samplers and our sample containers. This essentially provided a closed system of liquid handling which greatly simplified contamination control. The concentrations of acidic gases in our samples were quantified using

a custom built dual ion chromatography system equipped with a computer interface for data acquisition. The system was composed primarily of Dionex components with the detectors and flow system thermostated to 40°C. Eluants were constantly purged with He gas. Nitric acid was measured using a fast anion column while the carboxylic acids were determined using an AS4 column. Concentrator columns and electronic suppression was used in both chromatography systems. Calibration curves generated on the ground and in the air agreed within $\pm 2\%$. We thus were able to determine atmospheric mixing ratios of acidic gases in near-real time.

In addition to data for acidic gases, we present selected information on several important trace gases including ozone (O_3), carbon monoxide (CO), ethyne (C_2H_2), perchloroethylene (C_2Cl_4), and peroxyacetylnitrate (PAN). Aerosol NO_3^- was measured on bulk filter samples collected with a forward facing isokinetic probe housed in a shroud to ensure isoaxial flow [Dibb *et al.*, 1996a]. Ninety millimeter diameter 2 μm pore-sized Zefluor teflon filters were used as the collection substrate. Specific details regarding the measurement of various other species used in this paper are presented in companion papers [Blake *et al.*, this issue; Dibb *et al.*, this issue; Gregory *et al.*, this issue; Vay *et al.*, this issue]. The measurements of these species were averaged to provide mean values that corresponded directly to the acidic gas sampling times. This merged data product was generated at Harvard University, and it is used exclusively in this paper.

3. Results

In the data presented in this paper, obvious stratospherically impacted values have been removed based on coincident measurements of O_3 , CO, dew point, and selected hydrocarbons and halocarbons. This amounted to removing a total of about 25 data points obtained on three different flights.

The large-scale latitudinal distribution of acidic gases over the central and South Pacific basins is presented in Figure 1. It is evident from these distributions that numerous air parcels were encountered between 15° and 60°S latitude which contained large mixing ratios of acidic gases. The northern border of the impacted Pacific troposphere appears to be controlled by the presence and location of the South Pacific Convergence Zone (SPCZ) [Gregory *et al.*, this issue]. Nitric acid mixing ratios, for example, typically decreased by a factor of 2-5 in crossing the SPCZ region from south to north. This trend was also apparent in other trace gases such as CO, C_2H_2 , C_2H_6 , O_3 , and PAN [Gregory *et al.*, this issue]. Thus polluted air parcels appeared to be present south of the SPCZ with "clean" air north of it fed by an easterly flow regime along the southern edge of the ITCZ.

Mixing ratios of acidic gases over the South Pacific were generally less than 200 pptv but approached or exceeded 1000 pptv in some air parcels. These air parcels (i.e., plumes) were observed mainly between 2 and 7 km altitude (Figures 2a-2c). Because of the strong trade wind inversion over this region, the marine boundary layer exhibited very small mixing ratios of acidic gases. The inversion appeared to be a very effective barrier to downward mixing of acidic gases from aloft. Indeed, the most processed (i.e., aging and mixing influences) air parcels were sampled in the marine boundary layer. This feature of the data is illustrated using the ratio C_2H_2/CO which had a median value of 0.6 below 1 km altitude but showed significantly larger values in the rest of the tropospheric column (Figure 3). Values of this ratio less than 1.0 are typical of photochemically aged and well mixed (diluted) air parcels [Smyth *et al.*, 1998; Talbot *et al.*, 1997b].

As an example of the detailed vertical structure over the South Pacific selected data from a slow spiral (80 m min⁻¹) conducted east of Fiji is shown in Figure 4. An apparent combustion plume was sampled near 5 km, with corresponding large increases in HNO_3 , C_2H_2 , and C_2H_2/CO but not C_2Cl_4 . Notice the very rapid vertical changes in the mixing ratios and generally good correspondence between HNO_3 and C_2H_2 . While the plumes with large mixing ratios of many trace gases clearly stand out, the PEM-Tropics data in general support the idea that much of the tropospheric column from 2-10 km altitude was fumigated with varying degrees of combustion emissions. The smooth shape of the vertical distribution of C_2Cl_4 is typical of what was observed over the South Pacific (Figure 1), and it suggests minimal influence on the chemistry from industrial emissions. The distribution of CH_3CCl_3 and other halocarbons also supports this ascertain (N. Blake, personal communication, 1998).

To provide a detailed description of the distribution of acidic gases over the central and South Pacific basins, this information is presented in a regional summary format (Table 1) consistent with that used in companion papers [Gregory *et al.*, this issue; Dibb *et al.*, this issue]. Information on the distribution of many other trace gases can be found in these papers, so it is not duplicated here. The regional breakdown was developed to provide data summaries that correspond to logical latitudinal and longitudinal areas (e.g., the ITCZ, and the eastern, central, and western Pacific basins). In some regions the sampling was quite sparse, so interregional comparisons need to be conducted with caution. On the basis of the vertical measurement density of acidic gases, the data were broken into four altitude bins: (1) the marine boundary layer (<1 km), (2) the transition or cloud layer (1-2 km), (3) the middle (2-8 km), and (4) upper (8-12 km) troposphere.

As shown in the large-scale vertical distributions (Table 1 and Figure 2), the smallest mixing ratios of acidic gases were found in the marine boundary layer. Here median mixing ratios were 14 pptv for HNO_3 , 19 pptv for $HCOOH$, and 18 pptv for CH_3COOH . The very small mixing ratios of HNO_3 are consistent with the observed NO_x values of only a few or sub (i.e., <1) pptv in the boundary layer (Georgia Institute of Technology NO_x data are available from the Distributed Active Archive Center (DAAC) at NASA Langley Research Center, Hampton, Virginia). There was no significant regional difference in the mixing ratio of HNO_3 in the marine boundary layer, but the carboxylic acids exhibited values 2-3 times larger in the central Pacific region. In the middle troposphere the mixing ratios of acidic gases showed the largest values in the western and central regions. This is consistent with the generally westerly flow of air at these altitudes over the South Pacific basin, implying that the least processed air parcels would be found in these regions [Fuelberg *et al.*, this issue]. Most of the plumes that we sampled were, in fact, encountered over the western and central Pacific areas. The eastern Pacific regions were dominated by relatively "clean" air parcels. This longitudinal difference seemingly reflects chemical and physical losses of acidic gases as air parcels transverse the Pacific basin in a westerly flow regime.

The mixing ratios of the carboxylic acids $HCOOH$ and CH_3COOH are generally found to be highly correlated in the gas and liquid phases in the troposphere [Keene and Galloway, 1986]. Over continental areas the ratio $HCOOH/CH_3COOH$ usually has a value near 2.0 with a correlation coefficient between these two species near 0.9 [Keene and Galloway, 1986; Talbot *et al.*, 1988]. Although we observed linear correlations between $HCOOH$ and CH_3COOH over the Pacific basin (Figures 5a and 5b), they were less robust than what we observed during the Pacific Exploratory Mission-West Phases A and B (PEM-West A and B) [Talbot *et al.*,




Figure 1. Latitudinal distribution of acidic gases at altitudes of 2–12 km over the central and southern Pacific basins.

1997a]. In some of the plumes, HCOOH was highly enhanced with regard to CH_3COOH , and the ratio $\text{HCOOH}/\text{CH}_3\text{COOH}$ had values as large as 5.0 (plume median equal to 1.6). This suggests the possibility of substantial photochemical production of HCOOH compared to CH_3COOH (or more efficient loss of CH_3COOH) in some of the plumes that we sampled over the South Pacific. This point is further explored in later sections of this paper.

4. Discussion

4.1. Altitude Range of 2–12 km

The large-scale impact of pollution over much of the western and central Pacific basins is a significant feature of the PEM-Tropics

data set. Backward trajectories indicate that many of the air parcels we sampled had not been over continental areas for 10–20 days [Fuelberg *et al.*, this issue]. This is consistent with the chemical measurements which suggest that the air parcels were photochemically aged and physically processed for a week or two since the last injection of combustion emissions. Many of the trajectories follow a path that implies that the last continental areas that the air parcels passed over were Brazil and Africa. Since biomass burning occurs on both of these continental areas during austral spring [Cahoon *et al.*, 1992], this is likely to be a major source of combustion emissions over the Pacific basin at this time of year.

Methyl chloride is a reasonably good chemical tracer of biomass burning emissions [Blake *et al.*, 1996]. The relationship between




Figure 2. Vertical distribution of acidic gases over the central and southern Pacific basins. These distributions show that plume encounters with enhanced mixing ratios of acidic gases commonly occurred in the 3–7 km altitude region.

the mixing ratios of CH_3Cl and acidic gases is depicted in Figure 6. These plots indicate a general relationship between acidic gases and CH_3Cl ($r^2 = 0.4$). The enhancements of CH_3Cl in the plumes are small due to the significant dilution these well aged air parcels have undergone. Plots of C_2H_2 and C_2H_6 versus CH_3Cl (not shown) show similar relationships to those in Figure 6, again reflecting the substantial processing of the air parcels over the Indian and Pacific basins.

One feature of the plumes is the absence of enhancements in aerosol or aerosol associated species [Dibb *et al.*, this issue], even for ammonium which is released in large quantities from biomass combustion [Lobert *et al.*, 1991]. This indicates that the air parcels over the Pacific basin have been effectively scavenged by clouds and precipitation. It also suggests, since most acidic gases are highly water-soluble, that their large mixing ratios in some of the

plumes may be due to photochemical production since the last scavenging event. Evidence for a photochemical source of acidic gases is provided in Figures 7 and 8, where the relationships between these species and O_3 and PAN are presented. As with CH_3Cl , the trends are only general (r^2 near 0.4) but suggestive of photochemical production of acidic gases. The break in the relationships at <5 pptv of PAN is presumably due to thermal decomposition of PAN to NO_x at lower altitudes [Roberts, 1995]. The data corresponding to <5 pptv PAN was obtained in the 2–4 km altitude band where air temperatures were typically 280–285°K.

Plotting an individual species as a function of the ratio $\text{C}_2\text{H}_2/\text{CO}$ gives insight on the effect of air parcel processing on its mixing ratio. These relationships for acidic gases are shown in Figure 9. It is evident that the relationship is much tighter for the carboxylic acids compared to HNO_3 , but it is unclear as to why this is the case.




Figure 3. Vertical distribution of the ratio C_2H_2/CO over the central and southern Pacific basins.

Clearly, the largest mixing ratios of acidic gases were contained in the least processed air parcels ($C_2H_2/CO > 1$). On the basis of the correlations shown in this paper, it follows that these same air parcels also contained the largest mixing ratios of CH_3Cl , O_3 , and PAN. It appears that the chemical composition of these air parcels reflects photochemical activity of biomass burning emissions aged

over a minimum of a one week time frame. This is based largely on the absence of reactive hydrocarbons (i.e., C_4 and higher) in these air parcels. It appears that mixing processes (i.e., dilution) are responsible for much of the variation in individual species mixing ratios and inter-relationships between various compounds. Thus, air parcels can be quite photochemically aged with significant mixing ratios of secondary species but still appear much younger (e.g., $C_2H_2/CO > 1$) due to less mixing with background air.

To examine the potential combustion source of HNO_3 , only mixing ratios greater than 100 pptv are plotted versus CO and C_2H_2 (Figure 10). Only in a few plumes does there appear to be a direct relationship between HNO_3 and CO or C_2H_2 . The largest mixing ratios of HNO_3 occurred at relatively low values of CO and C_2H_2 and correspond to a C_2H_2/CO ratio near 1 (Figure 9). In general, there was very substantial amounts of HNO_3 in air parcels with CO of 50–100 ppbv and $C_2H_2 < 150$ pptv. Together these results point to significant photochemical production of HNO_3 (since the last scavenging event) during long-range transport of air parcels over the South Pacific. Similar arguments can be made for photochemical generation of carboxylic acids in these air parcels, especially $HCOOH$. Previous measurements of the ratio $HCOOH/CH_3COOH$ in biomass burning plumes transported long distances in the middle troposphere show values of 1.5–3 [Helas *et al.*, 1992; Lefer *et al.*, 1994; Dibb *et al.*, 1996b].

Additional modeling studies are needed to enhance our understanding of photochemical processes occurring within plumes over the South Pacific basin. Limited insight as to whether we observed loss of CH_3COOH in these plumes can be gleaned by examination of the CH_3COOH and CH_3OOH data. Plotting various subsets of these data obtained over the South Pacific (not shown) revealed no correspondence between the two species, as would be expected if CH_3COOH were a significant decomposition source of CH_3OOH .




Figure 4. Vertical distribution of selected trace gases during a slow spiral conducted during mission 17 just east of Fiji.

Table 1. Regional Summary of Acidic Gases Over the Central and Pacific Basins

Altitude, km	HNO ₃	HCOOH, pptv	CH ₃ COOH	N
<i>15°–45°N, 120°–170°W</i>				
< 1	13 ± 8.5 (10)	45 ± 18 (50)	55 ± 23 (60)	17
1–2	46 ± 10 (50)	39 ± 30 (30)	44 ± 35 (36)	6
2–8	87 ± 43 (83)	72 ± 49 (55)	72 ± 45 (59)	42
8–12	63 ± 53 (42)	55 ± 26 (52)	54 ± 30 (55)	29
<i>0°–15°N, 120°–170°W</i>				
< 1	20 ± 11 (15)	35 ± 30 (27)	34 ± 23 (20)	19
1–2	52 ± 14 (49)	33 ± 24 (18)	36 ± 26 (22)	10
2–8	67 ± 35 (51)	31 ± 14 (33)	33 ± 14 (35)	31
8–12	150 ± 107 (107)	27 ± 26 (14)	23 ± 19 (13)	16
<i>0°–35°S, 120°–170°W</i>				
< 1	18 ± 10 (17)	66 ± 216 (23)	84 ± 293 (27)	40
1–2	32 ± 14 (30)	44 ± 20 (43)	43 ± 345 (39)	21
2–8	139 ± 145 (84)	124 ± 181 (49)	84 ± 81 (54)	192
8–12	63 ± 63 (37)	61 ± 54 (46)	56 ± 46 (41)	122
<i>0°–35°S, > 170°W</i>				
< 1	15 ± 7.7 (13)	18 ± 7.4 (17)	17 ± 8.1 (15)	12
1–2	40 ± 3.0 (40)	50 ± 5.0 (50)	46 ± 8.5 (46)	2
2–8	160 ± 124 (97)	71 ± 54 (47)	84 ± 70 (53)	46
8–12	100 ± 61 (81)	66 ± 32 (52)	92 ± 27 (94)	19
<i>0°–35°S, 80°–120°W</i>				
< 1	13 ± 7.1 (14)	18 ± 5.9 (18)	16 ± 13 (18)	27
1–2	28 ± 8.2 (30)	33 ± 8.5 (32)	67 ± 47 (47)	5
2–8	43 ± 37 (31)	43 ± 28 (36)	52 ± 43 (36)	50
8–12	45 ± 30 (45)	54 ± 41 (39)	42 ± 26 (43)	28
<i>35°–72°S, > 170°W</i>				
< 1	14 ± 4.6 (14)	11 ± 4.1 (10)	13 ± 3.9 (12)	23
1–2	25 ± 9.9 (28)	38 ± 26 (33)	29 ± 16 (26)	13
2–8	180 ± 262 (80)	162 ± 176 (90)	114 ± 12 (57)	70
8–12	54 ± 71 (25)	54 ± 39 (40)	45 ± 25 (37)	23
<i>35°–72°S, 80°–120°W</i>				
< 1	12 ± 2.3 (11)	16 ± 4.4 (18)	18 ± 4.6 (19)	7
1–2	31 ± NA (31)	22 ± NA (22)	20 ± NA (20)	1
2–8	26 ± 16 (23)	35 ± 19 (36)	33 ± 18 (26)	16
8–12	38 ± 60 (9.0)	18 ± 5.0 (19)	16 ± 4.8 (17)	5

Values are stated as mean ± one standard deviation (median). N represents the number of data in altitude bin. NA means not applicable.




Figure 5. Relationship between mixing ratios of HCOOH and CH₃COOH over the central and southern Pacific basins; (a) <2 km and (b) 2–12 km altitude.




Figure 6. Relationships between mixing ratios of acidic gases and CH_3Cl in the altitude range 2–12 km. The r^2 values for these correlations were ≈ 0.35 . Particularly for HCOOH and CH_3COOH , there is a general trend of enhanced mixing ratios at the largest values of CH_3Cl . These correlations potentially indicate an important biomass burning source for acidic gases over the South Pacific basin. Note that the abscissas are also a logarithmic scale, ranging from 500 to 650 pptv.

[Madronich and Calvert, 1990]. This result appears to support a significant photochemical source of HCOOH rather than a predominance of decomposition of CH_3COOH in aged biomass burning plumes over the South Pacific. We can not rule out, however, some photochemical production of CH_3COOH as well. Because of potentially complex (and unknown) chemistry in these biomass

burning plumes, there is clearly much uncertainty surrounding the production and decomposition of carboxylic acids in such air parcels.

4.2 Altitude Range of 1–2 km

The altitude band <2 km was broken into the marine boundary layer (<1 km) and the transition or cloud layer from 1–2 km. The mixing ratio of acidic gases in these two layers are shown as a function of latitude in Figures 11a–11c. Nitric acid mixing ratios were smaller at <1 km compared to 1–2 km altitude, except for the most southerly data where they were about equal. There does not




Figure 7. Relationships between mixing ratios of acidic gases and O_3 in the altitude range 2–12 km. The r^2 values for these correlations were ≈ 0.40 . These general correlations potentially indicate a photochemical source for HCOOH and CH_3COOH .




Figure 8. Relationships between mixing ratios of acidic gases and PAN in the altitude range 2–12 km. The r^2 values for these correlations were ≈ 0.40 . These general correlations potentially indicate a photochemical source for HCOOH and CH_3COOH . Mixing ratios of PAN below ≈ 5 pptv were observed in the altitude range 2–4 km, where thermal decomposition of PAN was apparently significant.

appear to be any systematic variation of HNO_3 mixing ratios at <1 km altitude with latitude. Although the data are somewhat scattered, HNO_3 mixing ratios appear to increase in the transition layer going from south to north latitude. This apparent trend is driven to a large extent by the low values near 60°S . At midlatitudes and in the tropics there was 2–3 times more HNO_3 in the transition layer than at <1 km altitude. This observation could be related to evaporation of cloud droplets releasing HNO_3 to the gas phase in the transition layer. This process would be most active near the ITCZ, which is where the largest mixing ratios of HNO_3 were observed at this

altitude. In both layers, aerosol NO_3^- mixing ratios were about a factor of 2 greater than those of HNO_3 [Dibb *et al.*, this issue], presumably due to uptake of HNO_3 onto sea-salt particles in the marine boundary layer [Huebert, 1980] and possibly production of aerosol- NO_3^- from cloud processing in the transition layer.

The mixing ratios of carboxylic acids in the marine boundary layer over the South Pacific were about an order of magnitude less than those previously determined from shipboard sampling in the central North Pacific region [Arlander *et al.*, 1990]. This probably is due to the remoteness of the South Pacific basin from continental areas and restricted downward mixing across the trade wind inversion. Formic and acetic acid did not exhibit a difference in their mixing ratios between the marine boundary and transition layers. In the marine boundary layer they had the largest mixing ratios north of the ITCZ. This may reflect the closer proximity of




Figure 9. Distribution of the mixing ratios of acidic gases as a function of the ratio $\text{C}_2\text{H}_2/\text{CO}$.




Figure 10. Mixing ratio of HNO₃ as a function of CO and C₂H₂ for HNO₃ > 100 pptv.




Figure 11. Latitudinal distribution of acidic gases in the marine boundary layer (<1 km) and the overlying transition layer (1-2 km). The solid lines represent a plot of the median mixing ratio value as a function of latitude.




Figure 11. (continued)

continental areas to this region leading to enhanced primary or secondary production of these species. One difference between the vertical distribution of HNO_3 and the carboxylic acids in the two surface layers is the similarity of the mixing ratios of HCOOH and CH_3COOH in these two altitude bands but not those of HNO_3 . This observation may be explained by more extensive uptake of HNO_3 onto sea salt aerosols compared to the carboxylic acids. Detailed gas phase, cloud droplet, and aerosol measurements over the remote oceans are needed to better understand this issue.

5. Conclusion

The distribution of acidic gases over the South Pacific basin in austral springtime appears to be strongly influenced by emissions from biomass burning, most likely occurring in Africa and Brazil. Owing to the generally westerly flow of air over this area in the middle and upper troposphere, elevated mixing ratios of acidic gases and the presence of pollution plumes were concentrated in the western and central South Pacific. The eastern Pacific basin was relatively "clean" presumably due to chemical and physical removal of these species during the long transit across the South Pacific.

The enhanced mixing ratios of acidic gases in pollution plumes were coincident with relatively low mixing ratios of the combustion tracers CO and C_2H_2 . This observation and their general correlations with O_3 and PAN suggest that the mixing ratios of acidic gases may have been sustained by photochemical production in the pollution plumes. Most likely this generation of acidic gases occurred after the last scavenging event the air parcels encountered since other soluble species such as aerosols were not enhanced in these same plumes. The PEM-Tropics data document the hemispheric-scale pollution of the southern troposphere by biomass burning in austral springtime. The impact of these emissions on the chemistry of the southern hemisphere troposphere appears to be much greater than previously recognized.

Acknowledgments: We honor the outstanding unselfish contributions of our colleague and friend John Bradshaw (deceased) to the overall success and accomplishments of the GTE/PEM-Tropics airborne expedition. Excellent support was provided by the ground and flight crews of the NASA Ames DC-8 aircraft. This research was supported by the NASA Global Tropospheric Chemistry program.

References

- Andreae, M. O., R. W. Talbot, T. W. Andreae, and R. C. Harriss, Formic and acetic acid over the central Amazon region, Brazil, 1, Dry season, *J. Geophys. Res.*, **93**, 1616-1624, 1988.
- Andreae, M. O., R. W. Talbot, H. Berresheim, and K. C. Beecher, Precipitation chemistry in central Amazonia, *J. Geophys. Res.*, **95**, 16,987-16,999, 1990.
- Arlander, D. W., D. R. Cronn, J. C. Farmer, F. A. Menzies, and H. H. Westberg, Gaseous oxygenated hydrocarbons in the remote marine troposphere, *J. Geophys. Res.*, **95**, 16,391-16,403, 1990.
- Blake, N. J., D. R. Blake, B. C. Sive, T.-Y. Chen, and F. S. Rowland, Biomass burning emissions and vertical distribution of atmospheric methyl halides and other reduced carbon gases in the South Atlantic region, *J. Geophys. Res.*, **101**, 24,151-24,164, 1996.
- Cahoon, D. R., Jr., B. J. Stocks, J. S. Levine, W. R. Cofer III, and K. P. O'Neill, Seasonal distribution of African savanna fires, *Nature*, **359**, 812-815, 1992.
- Chameides, W. L., and D. D. Davis, Aqueous-phase source of formic acid in clouds, *Nature*, **304**, 427-429, 1983.
- Dibb, J. E., R. W. Talbot, K. I. Klemm, G. L. Gregory, H. B. Singh, J. D. Bradshaw, and S. T. Sandholm, Asian influence over the western North Pacific during the fall season: Inferences from lead-210, soluble ionic species, and ozone, *J. Geophys. Res.*, **101**, 1779-1792, 1996a.
- Dibb, J. E., R. W. Talbot, S. I. Whitlow, M. C. Shiphany, J. Winterle, J. McConnell, and R. Bales, Biomass burning signatures in the atmosphere and snow at Summit, Greenland: An event on 5 August 1994, *Atmos. Environ.*, **30**, 553-561, 1996b.
- Dibb, J. E., R. W. Talbot, E. M. Scheuer, D. R. Blake, N. J. Blake, G. L. Gregory, G. W. Sachse, and D. C. Thornton, Aerosol chemical characteristics and distribution during the Pacific Exploratory Mission, Tropics, *J. Geophys. Res.*, this issue.
- Folkens, I. A., A. J. Weinheimer, B. A. Ridley, J. G. Walega, B. Anderson, J. E. Collins, G. Sachse, R. F. Pueschel, and D. R. Blake, O_3 , NO_x , and NO_2/NO in the upper troposphere of the equatorial Pacific, *J. Geophys. Res.*, **100**, 20,913-20,926, 1995.
- Fuehlberg, H. E., R. E. Newell, S. Longmore, Y. Zhu, D. J. Westberg, E. V. Browell, D. R. Blake, G. R. Gregory, and G. W. Sachse, A meteorological overview of the PEM-Tropics experiment, *J. Geophys. Res.*, this issue.
- Gregory, G. L., et al., Chemical characteristics of Pacific tropospheric air in the region of the ITCZ and SPCZ, *J. Geophys. Res.*, this issue.
- Hales, J. M., and M. T. Dana, Wet removal of sulphur compounds from the atmosphere, *Atmos. Environ.*, **12**, 389-400, 1979.
- Helas, G., H. Bingemer, and M. O. Andreae, Organic acids over equatorial Africa: Results from DECAFE 88, *J. Geophys. Res.*, **97**, 6187-6193, 1992.
- Hoell, J. M., Jr., et al., Pacific Exploratory Mission in the tropical Pacific: PEM-Tropics A, August-September 1996, *J. Geophys. Res.*, this issue.
- Huebert, B. J., Nitric acid and aerosol nitrate measurements in the equatorial Pacific region, *Geophys. Res. Lett.*, **7**, 325-328, 1980.
- Jacob, D. J., Chemistry of OH in remote clouds and its role in the production of formic acid and peroxyoxymonosulfate, *J. Geophys. Res.*, **91**, 9807-9826, 1986.
- Jacob, D. J., and S. C. Wofsy, Photochemical production of carboxylic acids in a remote continental atmosphere, in *Acid Deposition Processes at High Elevation Sites*, edited by M. H. Unsworth, pp. 73-92, D. Reidel, Norwell, Mass., 1988.
- Kawamura, K., L.-L. Ng, and I. R. Kaplan, Determination of organic acids (C_1 - C_{10}) in the atmosphere, motor exhausts, and engine oils, *Environ. Sci. Technol.*, **19**, 1082-1086, 1985.
- Keene, W. C., and J. N. Galloway, Considerations regarding sources for formic and acetic acids in the troposphere, *J. Geophys. Res.*, **91**, 14,466-14,474, 1986.
- Keene, W. C., J. N. Galloway, and J. D. Holden Jr., Measurements of weak organic acidity in precipitation from remote areas of the world, *J. Geophys. Res.*, **88**, 5122-5130, 1983.
- Lefer, B. L., et al., Enhancement of acidic gases in biomass burning impacted air masses over Canada, *J. Geophys. Res.*, **99**, 1721-1737, 1994.

- Lobert, J. M., D. H. Scharffe, W.-M. Hao, T. A. Kuhlbusch, R. Seuwen, P. Warneck, and P. J. Crutzen, Experimental evaluation of biomass burning emissions: Nitrogen and carbon containing compounds, in *Global Biomass Burning - Atmospheric, Climatic, and Biospheric Implications*, edited by J. S. Levine, 289-304, MIT Press, Cambridge, Mass., 1991.
- Logan, J. A., Nitrogen oxides in the troposphere: Global and regional budgets, *J. Geophys. Res.*, **88**, 10,785-10,807, 1983.
- Madronich, S., and J. G. Calvert, Permutation reactions of organic peroxy radicals in the troposphere, *J. Geophys. Res.*, **95**, 5697-5717, 1990.
- Madronich, S., R. B. Chatfield, J. G. Calvert, G. K. Moortgat, B. Veyret, and R. Lesclaux, A photochemical origin of acetic acid in the troposphere, *Geophys. Res. Lett.*, **17**, 2361-2364, 1990.
- Roberts, J. M., Reactive odd-nitrogen (NO_x) in the atmosphere, in *Composition, Chemistry, and Climate of the Atmosphere*, edited by H. B. Singh, pp. 176-215, Van Nostrand Reinhold, New York, 1995.
- Sanhueza, E., and M. O. Andreae, Emission of formic and acetic acids from tropical savanna soils, *Geophys. Res. Lett.*, **18**, 1707-1710, 1991.
- Smyth, S., et al., Comparison of the chemical signatures between air mass classes from the PEM experiments over the western Pacific, *J. Geophys. Res.*, in press, 1998.
- Talbot, R. W., K. M. Beecher, R. C. Harriss, and W. R. Cofer, III, Atmospheric geochemistry of formic and acetic acids at a midlatitude temperate site, *J. Geophys. Res.*, **93**, 1638-1652, 1988.
- Talbot, R. W., M. O. Andreae, H. Berresheim, D. J. Jacob, and K. M. Beecher, Sources and sinks of formic, acetic, and pyruvic acids over central Amazonia, 2, Wet season, *J. Geophys. Res.*, **95**, 16,799-16,811, 1990.
- Talbot, R. W., B. W. Mosher, B. G. Heikes, D. J. Jacob, J. W. Munger, B. C. Daube, W. C. Keene, J. R. Maben, and R. S. Artz, Carboxylic acids in the rural continental atmosphere over the eastern United States during the Shenandoah Cloud and Photochemistry Experiment, *J. Geophys. Res.*, **100**, 9335-9343, 1995.
- Talbot, R. W., et al., Large-scale distribution of tropospheric nitric, formic, and acetic acids over the western Pacific basin during wintertime, *J. Geophys. Res.*, **102**, 28,303-28,313, 1997a.
- Talbot, R. W., et al., Continental outflow over the western Pacific basin during February-March 1994: Results from PEM-West B, *J. Geophys. Res.*, **102**, 28,255-28,274, 1997b.
- Vay, S. A., B. E. Anderson, T. J. Conway, S. W. Sachse, J. E. Collins, d. R. Blake, and D. J. Westberg, Airborne observations of the tropospheric CO_2 distribution and its controlling factors over the South Pacific basin, *J. Geophys. Res.*, this issue.
- D. R. Blake and N. J. Blake, Department of Chemistry, University of California, Irvine, CA 92717.
- J. E. Dibb, E. M. Scheuer, and R. W. Talbot, Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, NH 03824. (e-mail: rwt@christa.unh.edu)
- G. L. Gregory and G. W. Sachse, NASA Langley Research Center, Hampton, VA.
- S. T. Sundholm, Georgia Institute of Technology, Atlanta, GA 30332.
- H. B. Singh, NASA Ames Research Center, Moffett Field, CA 94035.

(Received October 10, 1997; revised March 2, 1998; accepted March 10, 1998.)